

2020 Florida Youth Substance Abuse Survey

Big Bend Community Based Care Data Tables

Table 1. Major demographic characteristics of surveyed Big Bend Community Based Care and Florida Statewide youth, 2020

	Big Bend Community Based Care		Florida Statewide	
	N	%	N	%
Sex				
Female	5,592	49.3	25,191	48.4
Male	5,674	50.0	26,328	50.5
Race/Ethnic group				
African American	2,136	18.8	11,284	21.7
American Indian	330	2.9	491	0.9
Asian	193	1.7	671	1.3
Hispanic/Latino	671	5.9	11,091	21.3
Native Hawaiian/Pacific Islander	37	0.3	68	0.1
Other/Multiple	2,058	18.1	7,227	13.9
White, non-Hispanic	5,839	51.4	20,904	40.1
Age				
10	19	0.2	44	0.1
11	764	6.7	3,151	6.0
12	1,816	16.0	7,124	13.7
13	2,023	17.8	7,698	14.8
14	1,941	17.1	7,616	14.6
15	1,651	14.5	7,582	14.6
16	1,309	11.5	7,502	14.4
17	1,110	9.8	7,073	13.6
18	614	5.4	3,807	7.3
19 or older	84	0.7	394	0.8
Grade				
6th	2,109	18.6	7,718	14.8
7th	1,956	17.2	7,555	14.5
8th	2,099	18.5	7,632	14.6
9th	1,638	14.4	7,668	14.7
10th	1,383	12.2	7,481	14.4
11th	1,136	10.0	7,117	13.7
12th	1,029	9.1	6,923	13.3
Overall Middle School	6,164	54.3	22,904	44.0
Overall High School	5,186	45.7	29,189	56.0
Total	11,350	100.0	52,093	100.0

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported having used various drugs in their lifetimes, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	22.5	43.1	36.3	31.1	24.2	43.1	33.7	23.0	44.9	38.3	32.5	24.4	44.8	35.3
Blacking Out	--	14.9	15.9	13.8	--	14.7	--	--	13.8	14.2	13.3	--	13.4	--
Cigarettes	9.2	15.4	11.5	13.4	9.3	15.2	12.6	6.1	11.3	8.5	9.6	6.2	11.1	9.0
Vaping Nicotine	15.7	34.6	27.9	24.1	16.8	35.2	26.0	13.5	30.0	24.9	20.8	14.8	30.1	22.8
Vaping Marijuana	7.2	20.8	15.2	14.1	7.7	21.2	14.6	6.8	22.2	16.6	14.5	7.5	22.3	15.5
Marijuana or Hashish	8.8	29.8	21.2	19.3	10.0	30.3	20.3	8.2	29.2	21.0	19.2	9.3	29.2	20.1
Synthetic Marijuana	--	3.2	2.7	3.6	--	3.2	--	--	2.9	2.9	2.9	--	2.8	--
Inhalants	8.7	5.4	7.8	5.7	8.5	5.6	6.9	8.3	5.1	7.7	5.2	8.1	5.0	6.5
Club Drugs	0.9	2.8	1.9	2.0	0.9	2.7	1.9	0.7	2.1	1.3	1.8	0.7	2.1	1.5
LSD, PCP or Mushrooms	1.1	4.9	2.8	3.5	1.2	5.0	3.2	1.1	4.3	2.5	3.4	1.1	4.3	2.9
Methamphetamine	0.8	1.2	0.9	1.2	0.8	1.2	1.1	0.7	0.8	0.6	0.9	0.6	0.9	0.8
Cocaine or Crack Cocaine	0.8	2.3	1.4	1.8	0.8	2.2	1.6	0.9	1.7	1.3	1.4	0.8	1.6	1.3
Heroin	0.5	0.8	0.5	0.8	0.5	0.9	0.7	0.5	0.6	0.4	0.6	0.4	0.6	0.5
Prescription Depressants	2.7	6.5	5.4	4.0	3.0	6.6	4.8	2.2	4.9	4.2	3.2	2.4	4.9	3.7
Prescription Pain Relievers	3.4	4.9	4.8	3.6	3.5	4.9	4.2	2.9	3.2	3.7	2.5	2.9	3.1	3.1
Prescription Amphetamines	2.6	5.5	4.9	3.3	2.9	5.4	4.1	2.1	4.5	3.7	3.2	2.3	4.4	3.4
Over-the-Counter Drugs	3.5	5.2	4.1	4.5	3.8	5.0	4.4	3.0	4.3	3.9	3.5	3.0	4.3	3.7
Needle to Inject Illegal Drugs	--	0.8	0.5	1.0	--	0.7	--	--	0.7	0.6	0.9	--	0.8	--
Any illicit drug	19.9	35.7	30.3	26.5	20.9	36.2	28.5	18.7	35.7	31.0	25.6	19.7	35.5	28.3
Any illicit drug other than marijuana	14.4	17.1	16.8	14.6	14.4	17.4	15.9	13.2	15.2	15.9	12.7	13.2	15.0	14.3
Alcohol only	10.9	15.3	13.6	13.0	11.7	15.0	13.3	11.9	17.1	15.3	14.3	12.5	17.1	14.8
Alcohol or any illicit drug	30.7	51.0	43.9	39.4	32.4	51.0	41.8	30.4	52.4	46.1	39.6	32.0	52.2	42.8
Any illicit drug, but no alcohol	8.4	8.0	7.8	8.5	8.4	8.0	8.2	7.5	7.7	7.9	7.3	7.7	7.5	7.6

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. For "Blacking Out," respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 3. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported having used various drugs in the past 30 days, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	7.9	20.0	16.0	12.9	8.7	19.5	14.5	8.2	19.9	15.9	13.7	8.7	19.7	14.8
Binge Drinking	3.4	9.2	6.3	6.6	3.6	8.5	6.5	3.4	9.2	6.8	6.4	3.6	8.9	6.7
Cigarettes	1.8	3.7	2.1	3.5	2.0	3.4	2.8	1.1	2.4	1.4	2.1	1.1	2.3	1.8
Vaping Nicotine	5.7	19.0	14.0	11.9	6.7	18.7	13.0	5.8	15.6	12.3	10.4	6.5	15.6	11.4
Vaping Marijuana	3.0	9.6	6.4	6.9	3.6	9.3	6.6	3.0	10.6	7.3	7.3	3.4	10.6	7.3
Marijuana or Hashish	3.5	14.4	9.5	9.4	4.4	13.9	9.4	3.8	15.9	10.6	10.7	4.4	15.8	10.7
Synthetic Marijuana	--	0.9	0.6	1.2	--	1.1	--	--	0.9	0.8	1.0	--	0.9	--
Inhalants	2.9	1.6	2.7	1.6	2.7	1.8	2.2	2.8	1.2	2.4	1.4	2.6	1.2	1.9
Club Drugs	0.4	1.1	0.7	0.9	0.4	1.2	0.8	0.3	0.6	0.4	0.6	0.3	0.7	0.5
LSD, PCP or Mushrooms	0.4	1.6	0.8	1.3	0.4	1.7	1.0	0.4	1.3	0.7	1.2	0.4	1.4	0.9
Methamphetamine	0.3	0.8	0.4	0.7	0.3	0.9	0.6	0.4	0.4	0.3	0.5	0.3	0.5	0.4
Cocaine or Crack Cocaine	0.3	0.8	0.4	0.7	0.3	0.9	0.6	0.3	0.5	0.4	0.5	0.3	0.6	0.4
Heroin	0.2	0.4	0.2	0.4	0.2	0.4	0.3	0.2	0.3	0.2	0.3	0.2	0.3	0.2
Prescription Depressants	0.8	1.6	1.3	1.1	0.9	1.6	1.2	0.8	1.2	1.0	1.0	0.8	1.3	1.0
Prescription Pain Relievers	1.5	1.3	1.8	1.1	1.5	1.3	1.4	1.2	1.0	1.3	0.8	1.2	1.0	1.1
Prescription Amphetamines	0.9	1.5	1.5	0.9	1.2	1.3	1.2	0.8	1.3	1.1	1.1	1.0	1.2	1.1
Over-the-Counter Drugs	1.4	1.7	1.5	1.5	1.6	1.6	1.5	1.4	1.3	1.4	1.2	1.4	1.3	1.3
Any illicit drug	9.0	18.3	14.6	13.3	9.8	18.1	14.0	8.8	19.5	15.5	14.1	9.2	19.4	14.8
Any illicit drug other than marijuana	6.0	6.4	6.5	5.8	6.0	6.7	6.2	5.7	5.4	6.0	5.0	5.5	5.5	5.5
Alcohol only	4.5	9.6	8.6	6.0	4.9	9.2	7.3	4.9	10.0	8.6	7.0	5.1	9.9	7.7
Alcohol or any illicit drug	13.4	27.9	23.2	19.3	14.6	27.2	21.3	13.6	29.1	23.9	20.8	14.2	29.0	22.3
Any illicit drug, but no alcohol	5.6	8.0	7.2	6.5	6.0	7.8	6.9	5.5	9.4	8.1	7.2	5.6	9.5	7.7

Note: The first 17 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Big Bend Community Based Care, 2010 to 2020

	2010			2012			2014			2016			2018			2020		
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total
Alcohol	35.4	62.5	50.6	28.3	59.6	45.7	23.4	54.3	40.6	22.1	52.5	39.3	22.1	47.6	36.4	22.5	43.1	33.7
Blacking Out	--	--	--	--	--	--	--	20.7	--	--	19.0	--	--	16.4	--	--	14.9	--
Cigarettes	21.6	37.4	30.5	16.3	33.4	25.8	12.5	29.3	21.8	10.4	26.3	19.3	10.0	20.2	15.6	9.2	15.4	12.6
Vaping Nicotine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	15.7	34.6	26.0
Vaping Marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	7.2	20.8	14.6
Vaporizer / E-Cigarette	--	--	--	--	--	--	--	--	--	13.8	37.8	27.2	15.3	38.4	28.1	--	--	--
Marijuana or Hashish	11.2	34.8	24.5	8.0	33.9	22.5	8.1	32.5	21.6	7.1	32.0	21.0	7.8	29.9	20.0	8.8	29.8	20.3
Synthetic Marijuana	--	--	--	--	16.0	--	--	9.3	--	--	6.7	--	--	3.4	--	--	3.2	--
Inhalants	13.6	9.0	11.0	10.7	7.8	9.0	8.2	5.8	6.8	6.8	4.9	5.8	8.0	5.2	6.4	8.7	5.4	6.9
Club Drugs	1.7	4.9	3.5	1.1	4.0	2.7	0.8	4.2	2.7	0.9	3.5	2.4	0.6	2.4	1.6	0.9	2.8	1.9
LSD, PCP or Mushrooms	2.3	5.6	4.2	1.3	5.2	3.5	1.1	5.7	3.7	1.0	5.8	3.7	0.7	4.4	2.7	1.1	4.9	3.2
Methamphetamine	1.2	1.6	1.4	1.0	1.4	1.2	0.7	1.3	1.0	0.6	1.0	0.9	0.5	1.0	0.8	0.8	1.2	1.1
Cocaine or Crack	1.9	3.8	3.0	1.0	3.9	2.7	0.8	2.7	1.9	0.9	3.3	2.2	0.7	2.5	1.7	0.8	2.3	1.6
Heroin	1.0	1.2	1.1	0.7	0.9	0.8	0.5	0.9	0.7	0.4	0.4	0.4	0.3	0.5	0.4	0.5	0.8	0.7
Prescription Depressants	3.2	10.0	7.0	2.5	8.2	5.7	2.0	8.3	5.5	2.7	9.8	6.7	2.9	8.0	5.7	2.7	6.5	4.8
Prescription Pain Relievers	5.4	12.2	9.3	4.0	10.8	7.8	3.3	9.8	6.9	3.5	8.3	6.2	2.6	6.4	4.7	3.4	4.9	4.2
Prescription Amphetamines	2.4	9.6	6.5	2.0	8.1	5.4	1.3	7.7	4.9	1.8	7.4	4.9	1.5	5.7	3.8	2.6	5.5	4.1
Over-the-Counter Drugs	5.8	9.4	7.8	4.2	7.6	6.1	3.6	6.5	5.2	3.2	6.5	5.0	3.5	6.2	5.0	3.5	5.2	4.4
Needle to Inject Drugs	--	--	--	--	--	--	--	--	--	--	0.9	--	--	0.8	--	--	0.8	--
Any illicit drug	25.0	41.8	34.5	19.5	41.8	31.9	17.7	39.6	29.8	16.0	37.3	27.9	17.1	36.2	27.7	19.9	35.7	28.5
Any illicit drug other than marijuana	20.4	26.2	23.7	16.0	24.1	20.6	13.6	22.8	18.7	12.6	21.5	17.6	13.2	19.1	16.5	14.4	17.1	15.9
Alcohol only	17.7	25.1	21.8	15.2	22.5	19.3	12.7	20.3	16.9	12.4	20.4	16.9	11.4	18.1	15.2	10.9	15.3	13.3
Alcohol or any illicit drug	42.7	66.8	56.2	34.5	64.1	51.0	30.3	59.6	46.6	28.0	57.6	44.5	28.1	54.2	42.6	30.7	51.0	41.8
Any illicit drug, but no alcohol	7.4	4.4	5.7	6.4	4.6	5.4	7.0	5.4	6.1	6.5	5.3	5.8	6.3	6.7	6.5	8.4	8.0	8.2

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. For "Blacking Out," respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. In 2020, the vaping/e-cigarette item was replaced with separate questions about vaping nicotine and vaping marijuana. The symbol "--" indicates that data are not available.

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Big Bend Community Based Care, 2010 to 2020

	2010			2012			2014			2016			2018			2020		
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total
Alcohol	16.5	37.6	28.4	11.5	34.1	24.1	9.9	29.2	20.6	8.2	27.6	19.2	8.5	23.5	16.9	7.9	20.0	14.5
Binge Drinking	7.2	20.8	14.9	4.5	17.8	11.9	3.8	15.1	10.0	2.9	13.5	8.9	2.8	11.5	7.6	3.4	9.2	6.5
Cigarettes	6.1	15.5	11.4	4.4	12.9	9.1	3.2	9.0	6.4	2.4	7.5	5.2	1.7	5.6	3.9	1.8	3.7	2.8
Vaping Nicotine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.7	19.0	13.0
Vaping Marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	3.0	9.6	6.6
Vaporizer / E-Cigarette	--	--	--	--	--	--	--	--	--	4.7	14.9	10.4	6.0	21.9	14.8	--	--	--
Marijuana or Hashish	5.9	19.2	13.4	3.8	17.6	11.6	4.1	16.8	11.1	3.2	16.0	10.4	3.5	15.1	10.0	3.5	14.4	9.4
Synthetic Marijuana	--	--	--	--	4.8	--	--	1.6	--	--	1.3	--	--	1.2	--	--	0.9	--
Inhalants	5.6	2.4	3.8	3.9	2.0	2.8	2.8	1.5	2.1	2.3	1.1	1.6	3.0	1.3	2.1	2.9	1.6	2.2
Club Drugs	0.5	1.4	1.0	0.3	1.3	0.9	0.4	1.1	0.8	0.3	0.9	0.6	0.3	0.7	0.5	0.4	1.1	0.8
LSD, PCP or Mushrooms	0.9	1.3	1.1	0.5	1.4	1.0	0.5	1.5	1.1	0.3	1.4	0.9	0.4	1.3	0.9	0.4	1.6	1.0
Methamphetamine	0.7	0.6	0.7	0.3	0.6	0.5	0.4	0.5	0.5	0.3	0.4	0.4	0.2	0.4	0.3	0.3	0.8	0.6
Cocaine or Crack	0.8	0.9	0.9	0.4	1.0	0.7	0.4	0.9	0.7	0.3	1.0	0.7	0.3	0.7	0.5	0.3	0.8	0.6
Heroin	0.4	0.4	0.4	0.2	0.5	0.4	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.4	0.3
Prescription Depressants	1.5	3.4	2.6	0.9	3.2	2.2	0.7	2.9	1.9	1.0	3.8	2.6	0.9	2.4	1.7	0.8	1.6	1.2
Prescription Pain Relievers	2.9	5.1	4.1	1.6	3.7	2.8	1.6	3.7	2.8	1.5	3.0	2.4	0.9	1.9	1.5	1.5	1.3	1.4
Prescription Amphetamines	1.2	3.4	2.5	0.8	2.8	2.0	0.7	2.6	1.7	0.6	2.3	1.5	0.6	1.6	1.2	0.9	1.5	1.2
Over-the-Counter Drugs	2.9	3.6	3.3	1.7	2.9	2.4	1.9	3.0	2.5	1.5	2.2	1.9	1.5	2.2	1.9	1.4	1.7	1.5
Any illicit drug	13.4	23.9	19.4	9.0	22.9	16.8	8.6	21.6	15.8	7.3	19.3	14.0	7.8	18.4	13.7	9.0	18.3	14.0
Any illicit drug other than marijuana	10.3	11.8	11.2	7.0	10.9	9.2	6.2	10.0	8.3	5.4	8.5	7.1	5.7	7.3	6.6	6.0	6.4	6.2
Alcohol only	9.5	20.0	15.4	7.3	17.5	13.0	6.0	15.1	11.0	5.4	15.1	10.9	5.3	12.7	9.4	4.5	9.6	7.3
Alcohol or any illicit drug	22.6	43.5	34.4	16.1	40.0	29.4	14.4	36.3	26.5	12.4	34.1	24.5	12.9	30.3	22.6	13.4	27.9	21.3
Any illicit drug, but no alcohol	6.3	6.1	6.2	4.9	6.0	5.5	4.6	7.4	6.1	4.5	6.7	5.7	4.6	7.1	6.0	5.6	8.0	6.9

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. In 2020, the vaping/e-cigarette item was replaced with separate questions about vaping nicotine and vaping marijuana. The symbol "--" indicates that data are not available.

Table 6. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, and peer disapproval, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol	--	16.0	15.1	16.9	--	14.9	--	--	15.9	16.4	15.4	--	15.3	--
Drinking at least once a month	--	2.9	2.4	3.4	--	2.9	--	--	2.7	2.9	2.5	--	2.5	--
Smoke cigarettes	--	8.0	6.6	9.4	--	8.1	--	--	5.9	5.6	6.2	--	5.8	--
Smoke marijuana	--	8.2	6.9	9.4	--	8.3	--	--	8.2	7.6	8.8	--	7.9	--
Great risk of harm														
1 or 2 drinks nearly every day	47.8	43.1	47.3	43.4	47.3	43.5	45.2	49.1	43.1	48.4	43.1	48.3	43.7	45.7
5+ drinks once or twice a week	61.5	53.8	59.5	55.2	61.1	55.0	57.3	61.1	54.2	60.4	54.1	60.6	54.6	57.2
1+ packs of cigarettes per day	66.1	66.7	67.4	65.8	67.1	66.7	66.5	65.4	66.3	67.3	64.5	65.6	66.5	65.9
Vaping nicotine	50.9	40.0	47.2	43.0	50.2	41.1	44.9	50.4	40.5	46.7	43.1	49.3	40.7	44.8
Vaping marijuana	53.7	36.7	45.6	43.4	52.2	37.4	44.4	52.6	35.1	44.1	41.5	50.8	35.5	42.8
Marijuana once or twice a week	47.3	25.5	35.4	35.5	45.4	25.7	35.3	46.8	22.9	34.1	32.8	44.7	23.2	33.4
Try marijuana once or twice	33.0	17.2	24.2	24.6	31.7	17.0	24.3	32.3	16.0	22.7	23.6	30.7	16.2	23.1
Use prescription drugs	70.6	66.8	69.1	68.0	70.7	66.6	68.6	68.5	66.8	69.4	65.8	68.6	66.9	67.6
Think it would be wrong to...														
Smoke cigarettes	95.6	89.5	93.3	91.3	95.3	90.6	92.2	96.1	91.6	93.8	93.3	95.9	92.3	93.5
Drink alcohol regularly	89.0	70.5	78.6	79.0	87.6	71.6	78.8	88.7	70.3	77.4	79.2	87.6	70.7	78.3
Smoke marijuana	89.5	64.2	74.8	76.5	87.0	64.9	75.6	89.4	63.2	73.9	75.4	87.6	63.6	74.6
Vape nicotine	90.0	77.6	83.0	83.4	89.0	78.8	83.1	90.2	79.9	83.7	85.0	89.5	80.5	84.4
Vape marijuana	91.8	75.4	82.6	83.0	90.6	75.9	82.7	91.6	75.0	81.7	82.8	90.6	75.3	82.2
Use other illicit drugs	97.6	93.9	96.6	94.7	97.3	94.4	95.6	97.5	93.5	95.9	94.6	97.4	93.6	95.3
Friends think it would be wrong to...														
1 or 2 drinks nearly every day	91.6	81.6	87.6	84.5	90.8	82.0	86.1	91.3	82.8	87.6	85.7	90.9	82.7	86.6
Smoke tobacco	93.6	85.3	90.2	88.0	93.2	86.0	89.1	94.4	88.8	92.1	90.6	94.1	89.2	91.3
Smoke marijuana	88.4	63.2	74.0	75.2	86.3	63.6	74.6	87.9	61.8	73.0	73.6	86.2	61.5	73.3
Vape nicotine	87.2	69.2	76.2	78.4	85.7	69.8	77.3	87.8	72.7	78.3	80.3	86.4	72.8	79.3
Vape marijuana	90.4	70.3	79.2	79.6	88.6	70.8	79.3	90.0	70.7	78.9	79.5	88.6	70.6	79.2
Use prescription drugs	95.4	91.5	94.0	92.6	95.1	91.6	93.3	95.2	91.5	93.7	92.6	94.9	91.3	93.1

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

Table 7. Trends in early ATOD use and attitudes toward substance use for Big Bend Community Based Care, 2010 to 2020

	2010			2012			2014			2016			2018			2020		
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total
Early ATOD use																		
Try alcohol	--	26.5	--	--	26.0	--	--	21.9	--	--	20.8	--	--	17.9	--	--	16.0	--
Drink monthly	--	5.4	--	--	5.8	--	--	4.4	--	--	3.9	--	--	3.0	--	--	2.9	--
Smoke cigarettes	--	20.8	--	--	18.1	--	--	15.6	--	--	13.2	--	--	10.0	--	--	8.0	--
Smoke marijuana	--	12.4	--	--	11.8	--	--	11.4	--	--	11.4	--	--	9.8	--	--	8.2	--
Great risk of harm																		
Daily alcohol use	42.4	39.9	41.0	43.6	36.6	39.7	44.3	37.2	40.4	44.9	36.6	40.2	46.7	40.2	43.1	47.8	43.1	45.2
5+ drinks daily	--	--	--	--	--	--	57.8	49.4	53.1	59.2	48.3	53.1	61.0	51.8	55.9	61.5	53.8	57.3
Daily cigarette use	67.4	66.4	66.8	67.3	67.1	67.2	67.5	68.8	68.3	68.4	65.2	66.6	65.1	63.1	64.0	66.1	66.7	66.5
Vaping nicotine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	50.9	40.0	44.9
Vaping marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	53.7	36.7	44.4
Marijuana weekly*	--	--	--	--	--	--	55.3	29.4	40.9	53.9	26.4	38.5	49.8	24.5	35.8	47.3	25.5	35.3
Try marijuana	40.7	23.3	30.9	40.0	20.7	29.2	39.5	18.0	27.5	37.5	17.7	26.4	32.7	15.6	23.3	33.0	17.2	24.3
Prescription drugs	--	--	--	--	64.3	64.3	72.2	67.4	69.5	70.4	63.8	66.7	67.6	64.3	65.8	70.6	66.8	68.6
Think it wrong																		
Smoke cigarettes	88.1	70.2	78.1	92.3	77.0	83.8	93.6	80.7	86.5	95.0	82.5	88.0	94.6	85.1	89.3	95.6	89.5	92.2
Drink regularly	79.7	56.1	66.4	86.1	59.2	71.1	88.4	61.7	73.6	88.8	63.1	74.4	88.0	65.2	75.3	89.0	70.5	78.8
Smoke marijuana	88.0	67.1	76.2	90.8	68.5	78.4	89.5	65.4	76.2	90.2	63.7	75.3	88.5	60.9	73.1	89.5	64.2	75.6
Vape nicotine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	90.0	77.6	83.1
Vape marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	91.8	75.4	82.7
Other illicit drugs	96.0	92.6	94.1	97.4	93.9	95.4	97.9	93.2	95.3	97.9	92.5	94.9	97.2	94.0	95.4	97.6	93.9	95.6
Friends think it wrong																		
Daily alcohol use	--	--	--	--	--	--	89.6	73.3	80.5	90.6	75.9	82.4	90.9	78.0	83.7	91.6	81.6	86.1
Smoke tobacco	--	--	--	--	--	--	92.5	79.1	85.1	93.3	82.1	87.1	93.6	82.7	87.6	93.6	85.3	89.1
Smoke marijuana	--	--	--	--	--	--	88.3	62.3	73.9	88.9	62.1	73.9	87.3	58.4	71.3	88.4	63.2	74.6
Vape nicotine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	87.2	69.2	77.3
Vape marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	90.4	70.3	79.3
Prescription drugs	--	--	--	--	--	--	96.0	88.3	91.7	95.8	87.8	91.3	94.6	88.8	91.4	95.4	91.5	93.3

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

* In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among high school drinkers, usual source of alcohol within the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2020

	Big Bend Community Based Care							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Bought in a store	--	4.5	3.0	6.8	--	3.9	--	--	6.5	4.6	8.9	--	5.9	--
Bought in a restaurant, bar, or club	--	1.5	1.1	2.0	--	0.9	--	--	1.9	1.9	2.0	--	1.5	--
Bought at a public event	--	0.8	0.0	1.9	--	1.0	--	--	1.2	0.7	1.7	--	1.1	--
Someone bought it for me	--	18.4	18.0	19.0	--	18.0	--	--	13.8	12.9	15.0	--	13.5	--
Someone gave it to me	--	41.7	43.3	39.2	--	42.7	--	--	40.2	42.9	36.9	--	41.1	--
Took it from a store	--	2.8	0.8	5.8	--	2.9	--	--	1.6	0.8	2.5	--	1.7	--
Took it from a family member	--	14.1	16.4	10.4	--	14.7	--	--	15.8	17.6	13.7	--	16.3	--
Some other way	--	16.2	17.4	14.9	--	15.8	--	--	19.0	18.7	19.3	--	18.8	--

Note: The symbol "--" indicates that data are not available.

Table 9. Among high school drinkers, usual drinking location within the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2020

	Big Bend Community Based Care							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
My home	--	37.5	40.6	32.9	--	36.5	--	--	42.7	44.1	40.8	--	43.3	--
Another person's home	--	40.1	39.7	40.4	--	42.2	--	--	35.2	37.2	33.0	--	35.4	--
Car or other vehicle	--	2.5	2.6	2.5	--	2.5	--	--	2.4	1.9	3.0	--	2.5	--
Restaurant, bar, or club	--	3.7	4.1	3.2	--	2.7	--	--	3.0	3.5	2.6	--	2.2	--
Public place	--	4.0	2.5	6.2	--	3.3	--	--	3.8	2.5	5.5	--	4.2	--
Public event	--	0.7	0.9	0.6	--	0.5	--	--	1.6	1.3	1.9	--	1.4	--
School property	--	1.4	0.7	2.4	--	1.7	--	--	1.2	1.0	1.4	--	1.2	--
Some other place	--	10.1	8.9	12.0	--	10.5	--	--	10.2	8.7	11.8	--	9.9	--

Note: The symbol "--" indicates that data are not available.

Table 10. Among high school drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2020

	Big Bend Community Based Care						Florida Statewide							
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
1	--	31.2	38.5	21.5	--	31.2	--	--	31.9	34.7	28.6	--	32.9	--
2	--	20.6	21.4	19.7	--	21.9	--	--	21.8	24.0	19.2	--	21.1	--
3	--	14.0	12.1	16.7	--	14.9	--	--	16.1	15.5	16.7	--	16.2	--
4	--	9.4	10.1	8.2	--	9.5	--	--	9.5	9.5	9.3	--	9.2	--
5 or more	--	24.9	17.9	33.8	--	22.5	--	--	20.7	16.3	26.2	--	20.5	--

Note: The symbol "--" indicates that data are not available.

Table 11. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high before or during school in the past 12 months, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Drinking alcohol	3.4	6.6	6.0	4.4	3.5	6.8	5.2	4.0	6.1	5.9	4.4	4.2	6.1	5.2
Smoking marijuana	3.3	11.3	7.6	7.6	3.8	11.3	7.6	4.0	13.6	9.9	9.1	4.5	13.9	9.5
Using another drug to get high	2.3	4.2	3.6	3.0	2.5	4.0	3.3	2.6	4.2	3.8	3.2	2.8	4.2	3.5

Table 12. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Carrying a handgun	7.4	7.3	4.5	10.2	7.1	7.2	7.3	6.2	6.0	3.6	8.6	6.0	5.9	6.1
Selling drugs	1.5	4.9	2.5	4.1	1.5	5.1	3.3	1.6	4.5	2.3	4.2	1.6	4.5	3.3
Attempting to steal a vehicle	1.1	2.0	1.3	1.8	1.0	2.0	1.6	1.3	1.6	1.1	1.8	1.3	1.5	1.5
Being arrested	2.4	2.7	2.1	3.1	2.1	2.9	2.6	2.2	2.3	1.8	2.7	2.1	2.3	2.3
Taking a handgun to school	0.4	0.9	0.5	0.9	0.4	0.8	0.7	0.4	0.7	0.4	0.8	0.5	0.6	0.6
Getting suspended	14.0	9.2	9.3	13.7	13.2	10.3	11.4	13.7	10.0	9.2	14.1	13.0	10.3	11.6
Attacking someone with intent to harm	8.3	6.0	5.6	8.5	8.2	6.1	7.0	8.3	5.5	5.8	7.6	7.8	5.6	6.7

Table 13. Trends in delinquent behaviors for Big Bend Community Based Care, 2010 to 2020

	2010			2012			2014			2016			2018			2020		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Carrying a handgun	6.0	5.7	5.8	5.8	4.9	5.3	6.4	6.7	6.5	7.1	6.7	6.9	7.0	7.7	7.4	7.4	7.3	7.3
Selling drugs	3.5	9.1	6.7	2.0	8.2	5.4	2.0	6.7	4.6	1.5	6.3	4.2	1.5	5.2	3.5	1.5	4.9	3.3
Attempting to steal a vehicle	2.1	2.2	2.1	1.2	2.6	2.0	1.1	1.8	1.5	0.6	2.0	1.4	1.1	1.6	1.4	1.1	2.0	1.6
Being arrested	3.7	4.9	4.4	2.8	4.3	3.6	2.3	3.0	2.7	1.9	3.0	2.5	2.1	2.9	2.5	2.4	2.7	2.6
Taking a handgun to school	0.9	1.1	1.0	0.4	0.9	0.7	0.4	0.8	0.6	0.3	0.8	0.6	0.4	1.0	0.7	0.4	0.9	0.7
Getting suspended	15.7	13.1	14.2	13.0	11.0	11.9	11.8	9.3	10.4	10.9	9.3	10.1	10.1	7.8	8.8	14.0	9.2	11.4
Attacking someone with intent to harm	11.9	10.7	11.2	7.6	8.2	7.9	6.8	6.8	6.8	6.8	6.8	6.8	8.1	6.0	6.9	8.3	6.0	7.0

Table 14. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported symptoms of depression, 2020

	Big Bend Community Based Care							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Sometimes I think that life is not worth it	30.2	33.0	41.7	21.6	30.6	33.2	31.7	29.9	31.4	39.5	21.9	29.9	31.7	30.7
At times I think I am no good at all	44.1	47.3	57.3	34.4	44.8	47.7	45.9	44.7	44.6	55.8	33.5	44.8	45.2	44.7
All in all, I am inclined to think that I am a failure	26.8	31.0	36.7	21.2	27.1	31.8	29.1	28.8	28.6	36.6	20.8	28.5	29.0	28.7
In the past year, have you felt depressed or sad MOST days, even if you felt OK sometimes?	45.7	50.0	59.0	37.0	45.9	50.4	48.0	44.5	47.6	57.0	35.5	44.6	48.0	46.2

Table 15. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported involvement in bullying behavior, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Skipped school because of bullying	8.6	11.1	14.6	5.1	8.5	11.5	10.0	8.8	9.7	13.3	5.4	8.8	9.8	9.3
Was ever kicked or shoved	43.0	29.8	35.6	35.6	41.7	30.9	35.7	39.8	24.8	30.7	31.9	38.8	24.8	31.4
Was ever taunted or teased	67.2	55.5	65.1	56.5	66.3	56.5	60.8	64.1	51.4	61.4	52.6	63.5	51.9	57.0
Was ever a victim of cyber bullying	27.2	32.3	38.8	21.4	27.2	33.2	30.0	25.7	29.0	35.0	20.4	26.1	29.2	27.5
Ever physically bullied others	24.0	13.2	15.3	20.7	22.9	13.6	18.0	22.4	12.5	14.2	19.3	21.3	12.6	16.8
Ever verbally bullied others	35.5	26.4	28.3	32.5	34.6	27.4	30.5	33.4	25.8	26.8	31.3	32.9	26.1	29.1
Ever cyber bullied others	11.4	12.8	12.9	11.5	11.4	13.3	12.2	11.2	12.4	12.3	11.5	11.4	12.4	11.9

Table 16. Percentages of Big Bend Community Based Care and Florida Statewide high school youth who reported adverse childhood experiences (ACEs), 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Emotional abuse	--	15.7	19.2	11.9	--	15.0	--	--	15.7	20.8	10.6	--	15.7	--
Physical abuse	--	10.5	11.1	9.7	--	10.4	--	--	10.5	11.9	9.1	--	10.5	--
Sexual abuse	--	6.0	9.5	2.2	--	6.2	--	--	5.5	8.7	2.2	--	5.7	--
Parents separated or divorced	--	42.7	45.0	40.4	--	42.4	--	--	40.1	43.1	37.1	--	40.6	--
Physical abuse in household	--	9.8	11.8	7.8	--	10.0	--	--	8.8	11.0	6.6	--	8.6	--
Substance abuse in household	--	30.7	33.0	28.3	--	31.2	--	--	26.8	30.3	23.1	--	26.5	--
Mental illness in household	--	34.7	41.7	27.4	--	34.6	--	--	30.5	38.0	22.8	--	30.6	--
Incarcerated household member	--	30.0	32.7	27.4	--	30.2	--	--	26.8	29.3	24.3	--	26.6	--
Emotional neglect	--	29.8	35.3	23.9	--	29.9	--	--	29.2	36.6	21.7	--	29.1	--
Physical neglect	--	7.9	9.0	6.8	--	8.0	--	--	7.4	8.7	6.1	--	7.2	--

Table 17. Number of adverse childhood experiences (ACEs) reported by Big Bend Community Based Care and Florida Statewide high school youth, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
No ACEs reported	--	30.4	26.2	34.7	--	30.3	--	--	32.1	25.8	38.4	--	31.7	--
1 ACE	--	19.5	17.8	21.4	--	19.4	--	--	21.0	20.0	22.1	--	21.4	--
2 ACEs	--	13.3	12.4	13.9	--	13.4	--	--	14.3	14.5	14.0	--	14.4	--
3 ACEs	--	12.3	14.3	10.4	--	12.3	--	--	10.9	12.1	9.8	--	10.8	--
4 ACEs	--	8.9	9.9	7.9	--	9.0	--	--	8.1	9.9	6.3	--	7.9	--
5 ACEs	--	6.6	7.9	5.2	--	6.6	--	--	5.7	6.9	4.5	--	5.8	--
6 ACEs	--	4.1	5.0	3.0	--	4.3	--	--	3.7	5.0	2.4	--	3.8	--
7 ACEs	--	2.4	3.1	1.7	--	2.2	--	--	2.1	2.8	1.2	--	2.0	--
8 ACEs	--	1.5	1.6	1.4	--	1.5	--	--	1.3	1.7	0.8	--	1.2	--
9 ACEs	--	0.7	1.0	0.3	--	0.7	--	--	0.6	0.9	0.2	--	0.6	--
10 ACEs	--	0.4	0.7	0.2	--	0.4	--	--	0.3	0.4	0.2	--	0.3	--

Note: Each respondent receives an ACEs score, which is the number of adverse childhood experiences he or she reported. This table shows the frequency distribution of the ACEs score across the sample. Percentages total to 100% down each column. Rounding can produce totals that do not equal 100%.

Table 18. Percentages of Big Bend Community Based Care and Florida Statewide youth who “agree” or “strongly agree” with statements indicating impulsiveness or a lack of self-control, 2020

	Big Bend Community Based Care							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
I often do what brings me pleasure now	27.1	25.2	27.8	24.1	26.8	26.6	26.1	32.0	28.8	30.9	29.4	31.5	29.2	30.2
I'm more concerned with the short run	24.7	20.3	23.6	20.8	24.2	20.7	22.3	28.1	23.1	26.8	23.8	27.6	23.1	25.3
Getting in trouble is exciting	26.6	28.8	28.1	27.2	26.9	29.5	27.8	27.5	28.2	29.1	26.6	27.6	28.8	27.9
Excitement is more important than security	22.7	21.5	20.9	23.1	22.8	21.5	22.0	25.1	23.9	23.3	25.5	25.0	24.3	24.4
People better stay away from me when I'm really angry	37.0	30.5	35.1	31.8	36.1	31.4	33.4	35.7	29.7	35.1	29.6	34.8	30.0	32.3
I get upset when I have a disagreement with someone	51.0	39.8	51.3	38.5	50.4	40.4	44.9	48.7	37.9	49.9	35.3	47.9	38.3	42.6

Note: The symbol “-” indicates that data are not available.

Table 19. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported participation in extracurricular activities, 2020

	Big Bend Community Based Care							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
School Sports	42.3	43.6	40.9	45.4	41.8	44.6	43.0	36.6	38.4	34.5	40.8	37.0	38.7	37.6
Organized Sports Outside of School	41.0	23.5	29.3	33.6	39.4	24.1	31.4	39.4	22.7	27.9	32.4	38.1	23.1	30.1
School Band	17.6	10.7	13.8	13.7	17.4	9.8	13.8	15.6	7.8	11.1	11.3	14.9	7.6	11.2
School Club(s)	25.3	35.0	38.7	22.9	25.7	35.4	30.7	22.7	31.5	35.3	20.2	23.0	31.8	27.6
Community Club(s)	9.0	12.3	13.4	8.4	8.8	12.3	10.8	8.6	12.8	13.8	8.3	8.7	12.5	11.0

Table 20. Percentages of high school students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Big Bend Community Based Care and Florida Statewide youth, 2020

	Big Bend Community Based Care							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
<i>Riding in a vehicle</i>														
Alcohol	--	15.7	16.6	14.7	--	15.8	--	--	14.7	16.2	13.3	--	14.8	--
Marijuana	--	22.4	24.5	20.5	--	22.5	--	--	22.1	24.0	20.2	--	21.8	--
<i>Driving a vehicle</i>														
Alcohol	--	4.5	4.1	4.8	--	4.5	--	--	4.1	3.6	4.6	--	3.8	--
Marijuana	--	8.8	8.7	8.9	--	8.6	--	--	9.0	8.2	9.8	--	8.8	--

Note: The symbol "--" indicates that data are not available.

Table 21. Trends for high school students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Big Bend Community Based Care, 2012 and 2020

	2010			2012			2014			2016			2018			2020		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
<i>Riding in a vehicle</i>																		
Alcohol	--	--	--	--	23.1	--	--	19.2	--	--	17.7	--	--	16.0	--	--	15.7	--
Marijuana	--	--	--	--	24.4	--	--	22.0	--	--	23.1	--	--	21.8	--	--	22.4	--
<i>Driving a vehicle</i>																		
Alcohol	--	--	--	--	9.7	--	--	7.4	--	--	6.6	--	--	4.9	--	--	4.5	--
Marijuana	--	--	--	--	11.4	--	--	11.2	--	--	10.6	--	--	9.6	--	--	8.8	--

Note: The symbol "--" indicates that data are not available.

Table 22. Protective factor prevalence rates for Big Bend Community Based Care, Florida Statewide youth and the national normative database, 2020

Domain	Scale	Big Bend Community Based Care		Florida Statewide		National Norms	
		Middle School	High School	Middle School	High School	Middle School	High School
Family	Family Opportunities for Prosocial Involvement	58	56	57	57	59	54
	Family Rewards for Prosocial Involvement	50	52	50	53	54	55
School	School Opportunities for Prosocial Involvement	52	62	51	60	57	60
	School Rewards for Prosocial Involvement	43	56	44	54	53	58
Peer and Individual	Religiosity	48	58	42	52	56	62
Average Prevalence Rate		50	57	49	55	56	58

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 23. Risk factor prevalence rates for Big Bend Community Based Care, Florida Statewide youth and the national normative database, 2020

Domain	Scale	Big Bend Community Based Care		Florida Statewide		National Norms	
		<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>
Community	Laws and Norms Favorable to Drug Use	40	30	41	32	42	42
	Perceived Availability of Drugs	34	20	34	20	45	45
	Perceived Availability of Handguns	28	36	23	31	25	42
Family	Poor Family Management	41	35	43	35	44	45
	Family Conflict	41	33	42	34	42	37
School	Poor Academic Performance	46	45	46	46	45	48
	Lack of Commitment to School	68	63	69	63	47	46
Peer and Individual	Favorable Attitudes toward Antisocial Behavior	48	37	49	38	40	46
	Favorable Attitudes toward ATOD Use	34	33	35	32	39	45
	Early Initiation of Drug Use	25	19	24	17	41	46
Average Prevalence Rate		40	35	41	35	41	44

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 24. Protective factor prevalence rate trends among middle school students for Big Bend Community Based Care, 2010 to 2020

Domain	Scale	Big Bend Community Based Care					
		2010	2012	2014	2016	2018	2020
Family	Family Opportunities for Prosocial Involvement	55	59	63	62	58	58
	Family Rewards for Prosocial Involvement	52	55	59	58	51	50
School	School Opportunities for Prosocial Involvement	47	54	57	57	56	52
	School Rewards for Prosocial Involvement	47	53	54	50	48	43
Peer and Individual	Religiosity	58	58	57	59	52	48
Average Prevalence Rate		52	56	58	57	53	50

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Risk factor prevalence rate trends among middle school students for Big Bend Community Based Care, 2010 to 2020

Domain	Scale	Big Bend Community Based Care					
		2010	2012	2014	2016	2018	2020
Community	Laws and Norms Favorable to Drug Use	46	38	38	37	38	40
	Perceived Availability of Drugs	48	39	38	38	33	34
	Perceived Availability of Handguns	31	29	31	31	29	28
Family	Poor Family Management	48	40	38	37	41	41
	Family Conflict	40	39	36	36	39	41
School	Poor Academic Performance	43	42	42	43	44	46
	Lack of Commitment to School	54	48	48	52	58	68
Peer and Individual	Favorable Attitudes toward Antisocial Behavior	48	40	37	39	42	48
	Favorable Attitudes toward ATOD Use	43	34	32	31	35	34
	Early Initiation of Drug Use	39	29	25	23	25	25
Average Prevalence Rate		44	38	37	37	38	40

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 26. Protective factor prevalence rate trends among high school students for Big Bend Community Based Care, 2010 to 2020

Domain	Scale	Big Bend Community Based Care					
		2010	2012	2014	2016	2018	2020
Family	Family Opportunities for Prosocial Involvement	56	57	59	61	59	56
	Family Rewards for Prosocial Involvement	55	55	58	56	53	52
School	School Opportunities for Prosocial Involvement	60	61	61	64	66	62
	School Rewards for Prosocial Involvement	61	61	61	59	59	56
Peer and Individual	Religiosity	67	67	66	64	58	58
Average Prevalence Rate		60	60	61	61	59	57

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 27. Risk factor prevalence rate trends among high school students for Big Bend Community Based Care, 2010 to 2020

Domain		Scale	Big Bend Community Based Care					
			2010	2012	2014	2016	2018	2020
Community	Laws and Norms Favorable to Drug Use		37	34	31	31	29	30
	Perceived Availability of Drugs		38	32	30	28	25	20
	Perceived Availability of Handguns		43	42	42	41	39	36
Family	Poor Family Management		44	40	38	36	34	35
	Family Conflict		34	33	31	33	33	33
School	Poor Academic Performance		45	44	40	44	42	45
	Lack of Commitment to School		51	46	49	53	55	63
Peer and Individual	Favorable Attitudes toward Antisocial Behavior		41	37	35	36	37	37
	Favorable Attitudes toward ATOD Use		42	39	39	38	38	33
	Early Initiation of Drug Use		34	32	28	26	21	19
Average Prevalence Rate			41	38	36	37	35	35

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.